



Real Biotech Corporation

13F.-2, No.33, Sec. 1, Minsheng Rd., Banqiao City, Taipei County 220, Taiwan, R. O. C.
Tel: +886 2 2950 9000 Fax: +886 2 2950 0505

www.real-biotech.com

HiYield™ Total RNA Mini Kit (Yeast)

Cat. No.:	YRY50	YRY100	YRY300
Product Name:	HiYield™ Total RNA Mini Kit (Yeast)		
Reactions:	50	100	300
Sample:	Up to 5×10^7 of a variety of yeast and fungus species		
Yield:	Up to 30µg		
Elution Volume:	50-100 µl		
Format:	Spin Column		
Operation:	Centrifuge		
Operation Time:	Within 20 Minutes		

Description

HiYield™ Total RNA Mini Kit (Yeast) is ideal for purification of total RNA from *Saccharomyces cerevisiae* and a variety of other yeast and fungus species in a simple spin column format. Sorbitol Buffer is included in the kit to reduce sample preparation time. Up to 30 µg of total RNA can be purified within 20 minutes without phenol/chloroform extraction or alcohol precipitation. Typical yield is around 20 µg for 5×10^7 of *Saccharomyces cerevisiae*. The convenient purification system removes contaminants well. Purified RNA is ready for use in RT-PCR, Northern Blotting, Primer Extension, mRNA Selection and cDNA Synthesis.

Features

1. Sorbitol Buffer is included in the kit to reduce sample preparation time.
2. Reliable isolation of ready-to-use RNA within 20 minutes.
3. Complete removal of all contaminants for sensitive downstream applications.

Contents

Item	YRY50	YRY100	YRY300
Sorbitol Buffer	45 ml	90 ml	225 ml
RB Buffer	30 ml	60 ml	130 ml
W1 Buffer	30 ml	50 ml	130 ml
Wash Buffer (concentrated)*	12.5 ml	25 ml	100 ml
RNase-free Water	6 ml	6 ml	30 ml
RB Column	50 pcs	100 pcs	300 pcs
2 ml Collection Tube	100 pcs	200 pcs	600 pcs

* Add absolute ethanol (see the bottle label for volume) to Wash Buffer then mix by shaking for a few seconds. Check the box on the bottle showing absolute ethanol is added. Be sure to close the bottle tightly after each use to avoid ethanol evaporation.



Real Biotech Corporation

13F.-2, No.33, Sec. 1, Minsheng Rd., Banqiao City, Taipei County 220, Taiwan, R. O. C.
Tel: +886 2 2950 9000 Fax: +886 2 2950 0505

www.real-biotech.com

Storage

HiYield™ Total RNA Mini Kit (Yeast) shall be shipped and stored dry at room temperature (15-25°C). With proper storage, HiYield™ Total RNA Mini Kit (Yeast) can be stored for up to 9 months without showing any deduction in performance and quality.

Applications

Purified RNA is ready for use in RT-PCR, Northern Blotting, Primer Extension, mRNA Selection and cDNA Synthesis.

Quality Control

The quality of HiYield™ Total RNA Mini Kit (Yeast) is tested on a lot-to-lot basis by isolation of total RNA from 5×10^7 of *Saccharomyces cerevisiae* harvested by centrifugation at 5,000 x g for 10 minutes. A 5 µl aliquot of purified RNA from a 50 µl eluate is analyzed by electrophoresis on a 0.8% agarose gel and the purified RNA with A260/A280 ratio 1.8 - 2.0 was quantified with a spectrophotometer.

Important Notes

Please read the entire notes prior to starting any of the protocol procedures.

Caution:

HiYield™ Total RNA Mini Kit (Yeast) contains irritants. During operation, always wear a lab coat, disposable gloves, protective goggles and (anti-fog) procedure mask.

Reagents to Be Supplied by User:

Lyticase or zymolase, absolute ethanol, ddH₂O (RNase-free and DNase-free) to prepare 70% ethanol, microcentrifuge tubes (RNase-free), pipette tips (RNase-free) and β-mercaptoethanol.

Things to Do before Starting:

Add absolute ethanol (see the bottle label for volume) to Wash Buffer then mix by shaking for a few seconds. Check the box on the bottle showing absolute ethanol is added. Be sure to close the bottle tightly after each use to avoid ethanol evaporation.

Steps to Prevent RNase Contamination:

1. During operation, always wear a lab coat, disposable gloves, protective goggles and (anti-fog) procedure mask.
2. Disposable plasticware and automatic pipettes should be sterile (RNase-free) and used only for RNA procedures.
3. Non-disposable glassware or plasticware should also be sterile (RNase-free).

DNA Removal Options:

For DNA-free RNA, perform optional steps listed in the protocol. Either follow the steps after RNA Binding, or steps after RNA Elution.

Protocol

Please read the entire important notes prior to starting.

<p>Step 1 Sample Preparation</p>	<ol style="list-style-type: none"> 1. Transfer fungus cells (up to 5×10^7) to a 1.5 ml microcentrifuge tube (RNase-free). 2. Harvest fungus cells by centrifugation for 10 minutes at 5,000 x g. 3. Discard the supernatant and resuspend the pellet in 600 μl of Sorbitol Buffer. 4. Add 2 μl of β-mercaptoethanol and 200 U of lyticase or zymolase. Then incubate at 30°C for 30 minutes. 5. Centrifuge the mixture for 10 minutes at 2,000 x g to harvest the spheroplast and then remove the supernatant.
<p>Step 2 Cell Lysis</p>	<ol style="list-style-type: none"> 1. Add 300 μl of RB Buffer and 3 μl β-mercaptoethanol to the sample lysate from Step 1 then vortex to mix. 2. Incubate the mixture at room temperature for 5 minutes. 3. Centrifuge at 14,000-16,000 xg for 2 minutes. 4. Transfer the supernatant to a new 1.5 ml microcentrifuge tube (RNase-free).
<p>Step 3 RNA Binding</p>	<ol style="list-style-type: none"> 1. Add 500 μl of 70% ethanol to the lysate and pipette immediately. 2. Place a RB Column in a 2 ml Collection Tube. 3. Transfer 500 μl of the mixture to the RB Column. 4. Centrifuge at 14,000-16,000 x g for 1 minute then discard the flow-through. 5. Transfer the remaining mixture to the same RB Column and centrifuge at 14,000-16,000 xg for 1 minute. Discard the flow-through and place the RB Column in a new 2 ml Collection Tube. <p><u>DNA Removal Option 1:</u> Add 100 μl of DNase I (2 KU/ml) mixed in a reaction buffer (50 mM Tris-HCl pH 7.5, 10 mM $MnCl_2$, 50 μg/ml BSA at 25°C) to the center of the RB Column matrix. Let stand for 10 minutes at room temperature and then proceed to Step 4 Wash.</p>
<p>Step 4 Wash</p>	<ol style="list-style-type: none"> 1. Add 400 μl of W1 Buffer to the RB Column then centrifuge at 16,000 x g for 1 min. 2. Discard the flow-through and place the RB Column back in the 2 ml Collection Tube. 3. Add 600 μl of Wash Buffer (ethanol added) into the RB Column. 4. Centrifuge at 14,000-16,000 x g for 1 minute. Discard the flow-through then place the RB Column back in the 2 ml Collection Tube. 5. Add 600 μl of Wash Buffer (ethanol added) into the RB Column. Centrifuge at 14,000-16,000 x g for 1 minute. Discard the flow-through then place the RB Column back in the 2 ml Collection Tube. 6. Centrifuge at 14-16,000 x g for 3 minutes to dry the column matrix.

<p>Step 5 RNA Elution</p>	<p>1. Place the dried RB Column in a clean 1.5 ml microcentrifuge tube (RNase-free).</p> <p>2. Add 50 μl of RNase-free Water into the CENTER of the column matrix. Let stand for at least 3 minutes to ensure the RNase-free Water is absorbed by the matrix.</p> <p>3. Centrifuge at 14,000-16,000 x g for 1 minute to elute the purified RNA.</p> <p><u>DNA Removal Option 2:</u></p> <p>Add 2 μl of DNase I (2 KU/ml) mixed in a reaction buffer (50 mM Tris-HCl pH 7.5, 10 mM MnCl₂, 50 μg/ml BSA at 25°C) to the final elution sample. Let stand for 10 minutes at room temperature.</p>
--------------------------------------	--

Troubleshooting

Problem	Possible Reasons/Solution
Clogged Column	<ol style="list-style-type: none"> 1. Reduce the amount of starting material or separate it into multiple tubes. 2. Centrifugation temperature must be between 20°C to 25°C. 3. Yeast cells were not completely homogenized. 4. Make sure lyticase or zymolase was added to Sorbitol Buffer immediately prior to use.
Residual Ethanol Contamination	<p>Following the wash step, dry the RB Column with additional centrifugation at 14,000-16,000 x g for 5 minutes.</p>
RNA Degradation	<ol style="list-style-type: none"> 1. The harvested sample should be stabilized immediately prior to use. 2. Disposable plasticware and automatic pipettes should be sterile (RNase-free) and used only for RNA procedures. 3. Non-disposable glassware or plasticware should also be sterile (RNase-free).